

VYBRANÁ LITERATURA

- Burba, G., Anderson, D. (2005): Introduction to the Eddy Covariance Method: general guidelines and conventional workflow. LI-COR Biosciences.
- Carboeurope (2000): Carboeurope Brochure 2000, Jena.
- Drexhage, M., Gruber, F. (1998): Architecture of the skeletal root system of 40-year-old *Picea abies* on strongly acidified soils in the Hartz Mountains (Germany). *Canadian Journal of Forest Research* **28**, 13–22.
- Dušek, J., Čížková, H., Czerný, R., Taufarová, K., Šmidová, M., Janouš, D. (2009): Influence of summer flood on the net ecosystem exchange of CO₂ in a temperate sedge-grass marsh. *Agricultural and Forest Meteorology* **149**, 1524–1530.
- Janssens, I. A., Freibauer, A., Ciais, P., et al. (2003): Europe's terrestrial biosphere absorbs 7–12 % of European anthropogenic CO₂ emissions. *Science* **300**, 1538–1542.
- Kankaala, P., Makela, S., et al. (2003): Midsummer spatial variation in CH₄ efflux from stands of littoral vegetation in a boreal meso-eutrophic lake. *Freshwater Biology* **48**, 1617–1629.
- Kellomäki, S., Oker-Blom, P., Kuuluvainen, T. (1984): The effect of crown and canopy structure on light interception and distribution in a tree stand. In *Crop Physiology of Forest Trees*. Tigerstedt, P. M. A., Puttonen, P., Koski, V. (Eds). University Press, Finland, pp. 107–115.
- Marek, M. V., Šprtová, M., Urban, O., Špunda, V., Kalina, J. (1999): Response of sun versus shade foliage photosynthesis to radiation in Norway spruce. *Phyton-Annales Rei Botanicae* **39**, 131–137.
- Mäkelä, A. (1997): A Carbon Balance Model of Growth and Self-Pruning in Trees Based on Structural Relationships. *Forest Science* **43**, 7–24.
- Pavelka, M., Acosta, M., Janouš, D. (2004): A new device for continuous CO₂ flux measurements in forest stand. *Ecology* **23** (2), 8–100.
- Pavelka, M., Acosta, M., Marek, M. V., Kutsch, W., Janouš, D. (2007): Dependency of the Q₁₀ values on the depth of soil temperature measuring point. *Plant and Soil* **292** (1–2), 171–179.
- Pokorný, R., Urban, O., Marek, M. V. (2004): Effect of Norway spruce planting density on shoot morphological parameters. *Biologia Plantarum* **48**, 137–139.
- Pokorný, R., Tomášková, I. (2007): Allometric relationships for surface area and dry mass of Norway spruce aboveground organs. *Journal of Forest Science* **53**, 548–554.
- Pokorný, R., Tomášková, I., Havránková, K. (2008): Temporal variation and efficiency of LAI in young mountain Norway spruce stand. *European Journal of Forest Research* **127**, 359–367.
- Prach, K. (1993): Vegetational changes in a wet meadow complex, South-Bohemia, Czech Republic. *Folia Geobotanica Phytotaxonomia* **28**, 1–13.
- Prach, K., Soukupová, L. (2002): Alterations in the Wet Meadows vegetation pattern. In *Freshwater Wetlands and Their Sustainable Future. A Case Study of the Třeboň Basin Biosphere Reserve*, Czech Republic. Květ, J., Jeník, J., Soukupová, L. (Eds). UNESCO Paris and Parthenon Publ. Boca Raton, pp. 243–254.
- Richardson, J. L., Vepraskas, M. J. (Eds) (2001): *Wetland Soils. Genesis, Hydrology, Landscapes and Classification*. Lewis Publishers, Boca Raton, 417 pp.
- Rinne, J., Ruita, T., Philatie, M., Aurela, M., Haapanala, S., Tuovinen, J. P., Tuitilla, E. S. (2007):

- Annual cycle of methane emission from a boreal fen measured by the eddy covariance technique. *Tellus* **59B**, 449–457.
- Vetter, M., Wirth, Ch., Böttcher, H., *et al.* (2005): Partitioning direct and indirect human-induced effects on carbon sequestration of managed coniferous forests using model simulations and forest inventories. *Global Change Biology* **11**, 810–827.
- Waring, R. H. (1987): Characteristics of trees predisposed to die: stress causes distinctive changes in photosynthate allocation. *Bio Science* **37**, 569–574.